

# **TCAD Avalanche Models**

**Group meeting  
19-05-2020**

# Preview

- **Wolff :**
  - First theoretical explanation of charge multiplication.
  - Solved quasi-Maxwellian equation with spherically symmetric distribution function.
  - Ionization coefficient valid for **high field** strengths :

$$\alpha(\epsilon) = a \exp(-b/\epsilon^2)$$


- **Shockley:**
  - A simple statistical model with three adjustable parameters
  - Spike distribution function.
  - Ionization coefficients valid for **low field** strengths :

$$\alpha(\epsilon) = (q\epsilon/rE_R) \exp(-E_i/qL_R\epsilon)$$

# Preview

- **Baraff :**
  - Distribution function corresponds to Wolff's spherically symmetric part and Shockley's spike with constant mean free path.
  - Similarities in slope to Wolff's theory for higher fields and Shockley's theory for lower fields.
- **Crowell & Sze :** modified Baraff's results
  - introduced parameters - average phonon energy lost per scattering and temperature.

$$\langle E_R \rangle = E_R \operatorname{tgh}(E_R/2kT)$$




- Later Keldysh confirmed Baraff's results.

# **TCAD Avalanche Models**

- **Van Overstraeten and de Man Model**
- **Okuto and Crowell Model**
- **Lackner Model**
- **UNIBO (University of Bologna) Model**
- **New UNIBO Model**

# Chynoweth Law

- Chynoweth's experimental ionization rates agreed well with measurement of ionization rates in gases :  $\alpha = a \exp(-b/\epsilon)$
- Observed linearity in  $\ln \alpha$  against  $E^{-1}$  graph over wide range of fields.
- Shockley and Van Overstraeten validated this law in the electric field range from  $1.75 \times 10^5 - 6 \times 10^5 \text{ Vcm}^{-1}$  (300K).



(A & B correspond to Chynoweth law)


# Van Overstraeten Model

- Used Chynoweth's law for fitting the data and extracting the model parameters :

$$\alpha(\epsilon_{ava}) = \gamma a \exp(-\gamma b/\epsilon_{ava});$$

$$\gamma = \tanh(\hbar\omega_{op}/2kT_0)/\tanh(\hbar\omega_{op}/2kT)$$

- Valid in the field range of  $1.75 \times 10^5 \leq \epsilon \leq 6 \times 10^5 \text{ Vcm}^{-1}$
- Different values of parameters **a** & **b** in low and high range of fields.
- Almost same value of **a** & **b** for 7 different diodes used - made good approximation of Chynoweth's law.



(Symbols represent particular diode)

# Van Overstraeten Model

Symbol	Parameter name	Electrons	Holes	Valid range of electric field	Unit
$a$	a (low)	$7.03 \times 10^5$	$1.582 \times 10^6$	$1.75 \times 10^5 \text{ Vcm}^{-1}$ to $E_0$	$\text{cm}^{-1}$
	a (high)	$7.03 \times 10^5$	$6.71 \times 10^5$	$E_0$ to $6 \times 10^5 \text{ Vcm}^{-1}$	
$b$	b (low)	$1.231 \times 10^6$	$2.036 \times 10^6$	$1.75 \times 10^5 \text{ Vcm}^{-1}$ to $E_0$	$\text{V/cm}$
	b (high)	$1.231 \times 10^6$	$1.693 \times 10^6$	$E_0$ to $6 \times 10^5 \text{ Vcm}^{-1}$	
$E_0$	E0	$4 \times 10^5$	$4 \times 10^5$		$\text{V/cm}$
$\hbar\omega_{\text{op}}$	hbarOmega	0.063	0.063		eV
$\lambda$	lambda	$62 \times 10^{-8}$	$45 \times 10^{-8}$		cm
$\beta$	beta (low)	0.678925	0.815009	$1.75 \times 10^5 \text{ Vcm}^{-1}$ to $E_0$	1
	beta (high)	0.678925	0.677706	$E_0$ to $6 \times 10^5 \text{ Vcm}^{-1}$	

# Okuto & Crowell Model

- Empirical model based on Baraff's theoretical model :

$$a(\epsilon_{ava}) = a [1+c(T-T_0)] \cdot \epsilon_{ava}^\gamma \exp(-(b[1+d(T-T_0)] / \epsilon_{ava})^\delta); \quad T_0=300K$$

- Energy conservation conditions applied.
- Applicable in the field range of  $10^5$ - $10^6$  Vcm $^{-1}$ .
- Applied pseudo-local approximation and predicted the existing data with good accuracy.
- “Exact” non-localized approximation considered boundary regions (p-i-n junction).
  - Zero ionization coefficient in boundary dark spaces (p-i & i-n)
  - Constant and close to the values of previous approximation in rest of the region.

# Okuto & Crowell Model

Symbol	Parameter name	Electrons	Holes	Unit
$a$	a	0.426	0.243	$V^{-1}$
$b$	b	$4.81 \times 10^5$	$6.53 \times 10^5$	$V/cm$
$c$	c	$3.05 \times 10^{-4}$	$5.35 \times 10^{-4}$	$K^{-1}$
$d$	d	$6.86 \times 10^{-4}$	$5.67 \times 10^{-4}$	$K^{-1}$
$\gamma$	gamma	1	1	1
$\delta$	delta	2	2	1
$\lambda$	lambda	$62 \times 10^{-8}$	$45 \times 10^{-8}$	cm
$\beta$	beta	0.265283	0.261395	1

# Lackner Model

- New theory on pseudo-local ionization probability model for field correction of Chynoweth law :

$$a_v(\epsilon_{ava}) = (\gamma a_v / Z) \exp(-\gamma b_v / \epsilon_{ava}); \text{ where } v = n, p$$

$$Z = 1 + (\gamma b_n / \epsilon_{ava}) \exp(-\gamma b_n / \epsilon_{ava}) + \gamma b_p / \epsilon_{ava} \exp(-\gamma b_p / \epsilon_{ava})$$

$$\gamma = \tanh(\hbar\omega_{op}/2kT_0) / \tanh(\hbar\omega_{op}/2kT)$$



- Introduced Z parameter in Chynoweth's law.
- Valid in the field range  $2 \times 10^5 - 6 \times 10^5 \text{ Vcm}^{-1}$ .
- Unlike Van Overstraeten and de Man model, same values of parameters **a** & **b** in low and high field regions.

# Lackner Model



Symbol	Parameter name	Electrons	Holes	Unit
$a$	a	$1.316 \times 10^6$	$1.818 \times 10^6$	$\text{cm}^{-1}$
$b$	b	$1.474 \times 10^6$	$2.036 \times 10^6$	V/cm
$\hbar\omega_{\text{op}}$	hbarOmega	0.063	0.063	eV
$\lambda$	lambda	$62 \times 10^{-8}$	$45 \times 10^{-8}$	cm
$\beta$	beta	0.812945	0.815009	I

# Lackner Model

- Field correction matched well with Chynoweth's law ( $Z=1$ ) for field values  $< 4 \times 10^5 \text{ Vcm}^{-1}$ .
- Deviation ( $Z \neq 1$ ) in higher range of fields ( $> 4 \times 10^5 \text{ Vcm}^{-1}$ ).
- No validation issue has been provided for field correction approximation.



# Comparison of Three Models



# UNIBO Model

- Compact model based on impact ionization data generated by Boltzmann solver HARM :

$$a(\epsilon_{ava}, T) = \epsilon_{ava} / (a(T) + b(T) \exp[d(T) / \epsilon_{ava} + c(T)])$$

- Ionization coefficient as a function of field and lattice temperature
- Developed for an extended temperature range (300K-675K) and low electric fields( $4 \times 10^4$  -  $5 \times 10^5$  Vcm $^{-1}$ ).
- Observed the contribution of non-equilibrium Auger generation at high temperatures - differentiate this model from other standard models.

# New UNIBO Model

- Modification of compact UNIBO model by extending the temperature range between 300-773K :

$$a(\epsilon_{ava}, T) = \epsilon_{ava} / (a(T) + b(T) \exp[d(T) / (\epsilon_{ava} + c(T))])$$

- Likewise UNIBO, New UNIBO model developed for low electric field range.
- Theoretically based on UNIBO theory
  - Different method of solving the parameters, so, resulted different values.

# Conclusions

- Van Overstraeten and Lackner model based on Chynoweth law, but used different parameters
  - reliable for low-field regions (Shockley's low field theory).
- Okuto and Crowell model valid for whole range of fields (Baraff's theory)..
- UNIBO and New UNIBO models showed their reliability for high temperature ranges and low electric fields(down to  $4 \times 10^4 \text{ Vcm}^{-1}$ ).
- All models except Okuto & Crowell typically used for self-heating power devices and ESD-protection structures.
- Our focus is on thin p-n junctions with moderate and high field regions - **Okuto and Crowell**.